# With information changing so quickly it's hard to keep up. Let us help.

# What masks should you wear and when?

Staff should follow the guidance of any facility they enter during the workday



### **Cloth Masks**

**Protects others**Reuse daily; Launder regularly

### Should be worn:

- ✓ Daily at work in a DHS office or congregate care setting
- At low-risk facilities where no one has been diagnosed with COVID-19 and social distancing is possible (ex: personal care homes)
- ✓ During home visits if you do not enter the home and no one in the home has a positive COVID-19 diagnosis/ symptoms
- ✓ In homeless shelters where no one has a positive COVID-19 diagnosis/ symptoms



## **Surgical Masks**

Protects self and others Wear up to 8 hours

#### Should be worn:

- When interacting with non-diagnosed individuals, including colleagues working in facilities with highrisk populations like congregate care settings
- ✓ In medium-risk facilities where there may have been positive COVID-19 cases and social distancing is difficult (ex: congregate care homes)
- ✓ During home visits where you do enter the home and no one in the home has a positive COVID-19 diagnosis/symptoms
- When working on isolation floors in the Safe-Haven Hotel where residents do not have a positive COVID-19 diagnosis, are not presumptive positive, or are not self-reporting symptoms
- ✓ In homeless shelters where daily mask laundering is not available, or where there may have been positive COVID-19 cases and social distancing is difficult



# N95/KN95 Masks

Highest protection for self and others

Wear up to 8 hours

### Should be worn:

- For any face-to-face interaction (clinical setting, home visit, or quarantine facility) with a client who has a positive COVID-19 diagnosis, is presumptive positive, or is self-reporting symptoms
- During any home situation and a family member has a positive COVID-19 diagnosis, is presumptive positive, or is self-reporting symptoms
- ✓ When working on floors in the Safe-Haven Hotel where residents have have a positive COVID-19 diagnosis, are presumptive positive, or are selfreporting symptoms

